

Analysis of the Thermodynamic Properties of a Binary Mixture of Pharmaceutical Residual Solvents

J. Ravi Kumar Reddy¹, K. Jyoshna Jayaraju², J. Niranjana³

Abstract

Deep examination on thermodynamic characteristics (Enthalpy, Entropy, Free energy) across the complete composition range from 298.15 K to 323.15 K at atmospheric pressure was used to study the molecular interaction of pharmaceutical residual solvents mixture (2-Butanol with m-Xylene). The solvation process, or the removal of molecules from their natural environment, is used to make sense of the data and provide context for the results. Then, the Redlich-Kister polynomial equation was used to fit the computed excess thermodynamic characteristics. The molecular interactions between 2-Butanol and m-Xylene are addressed in light of the computed excess characteristics.

Keywords: Residual solvent; Thermodynamic property; Binary mixture; 2-Butanol; m-Xylene

1. Introduction

In terms of output, the pharmaceutical sector is a major consumer of organic solvents [1-3]. Properties of binary liquid mixtures are of interest to pharmacists, medical scientists, condensed-matter theorists, experimental chemists, physicists, etc. since organic solvents are present throughout the pharmaceutical manufacturing process. Understanding the intermolecular forces responsible for the many phenomena seen in experiments is crucial for both fundamental and applied research. The physical, chemical, and transport characteristics of fluids are of interest since they are the most common form in which materials are handled in chemical process industries. Understanding the intermolecular interaction between the component molecules and processing product formation are both greatly aided by thermodynamic analysis of liquid mixtures consisting of polar and nonpolar components for a variety of industrial and technical applications. Testing for residual solvents has evolved into an integral aspect of drug quality assurance [4, 5].

Organic volatile molecules utilized or created in the manufacture of drug ingredients or excipients, or in the production of drug products, are referred to as residual solvents in the pharmaceutical industry. In terms of pharmacological significance, 2-butanol (or 2-Bu-OH) and xylene stand out among the residual solvents [6,7]. 2-Bu-OH is a polar chemical molecule that often appears as a racemic mixture of the two stereoisomers, (R)-(-)-2-Bu-OH and (S)-(+)-2-Bu-OH. This secondary alcohol is flammable, odorless, and colorless; it dissolves in water and is totally miscible with polar organic solvents like ethers and other alcohols [8, 9]. As a solvent, in the production of chemicals and pharmaceuticals, and in the field of medicine and dentistry [10, 11], xylene has several applications. The most prevalent form of xylene, known as m-xylene (or m-Xln), is a benzene derivative with two methyl groups substituted for the carbon atoms, where "m" stands for "meta," designating positions 1 and 3 for the two methyl substituents on the aromatic ring.

Professor & Principal¹, Associate Professor^{2,3}
Dept. of Pharmaceutics, Pharmacology

Mother Theresa Institute of Pharmaceutical Education and Research, Kurnool, Andhra Pradesh

Due to their widespread use, studies analyzing the nature of molecular interactions and exploring the physicochemical characteristics of such systems [12, 13, 14] have acquired a lot of traction in recent years. A thorough literature search revealed the lack of any comprehensive research focusing specifically on the thermodynamic parameters of the binary combination of 2-Bu-OH and m-Xln. In light of this relevance, it is of interest to research the thermodynamic characteristics in order to understand the interaction behavior in their binary mixes of various proportions with the variation of temperature. Thermodynamic characteristics and their excess thermodynamic properties have been computed from experimental measurements of density and viscosity. 298.15K to 323.15 K. The Redlich-Kister equation [15] was used to get the values for the excess characteristics, and the resulting enthalpy, entropy, and free energy of activation were then shown graphically.

2. Experimental details

2-Bu-OH and m-Xln used in this research study were purchased from Sigma Aldrich Chemicals Company. According to the manufacturer, the purities of these compounds were >99%. The water used in all experimental work was double distilled in quality. The binary mixtures of 2-Bu-OH and m-Xln were prepared by using an analytical electrical balance with a precision of $\pm 0.1 \mu\text{g}$ and later were converted to different composition of the mixture using dilution method. Special care was taken to prevent evaporation and the introduction of moisture into the experimental samples.

Density of all binary mixtures including pure solvents was measured using an oscillation densimeter (Anton Paar DSA 5000). To measure the viscosity Stabinger viscometer (svm-3000-stabinger-viscometer) was used. The temperature was previously set up by 298.15-323.15 K. The measurement was accomplished by the descending of the temperature in the viscometer. In both machinery processes the temperature was automatically controlled with an uncertainty of ± 0.01 K.

The plot of

$$\ln \left(\frac{\eta V_m}{hN} \right) \quad | \text{ ——— } |$$

vs. $1/T$ of eq. (3) gives a

3. Theoretical aspects

Calculation of different thermodynamic parameters for viscous flow:

Fluidity of liquid follows some mechanisms as liquid is considered as a combination of layers and so it flows as a rate process. To treat the viscosity as a rate process it is assumed that the motion of liquid layers involves the passage of a molecule from one equilibrium position to another in the same layer. To do so, it is necessary that a suitable „hole“ or site should be available, and the production of such a „hole“ requires the expenditure of energy, since work must be done in pushing back other molecules. The jump of the moving molecules from one equilibrium position to the next may thus be regarded as equivalent to the passage of the system over a potential barrier. Eyring and his co-workers using absolute reaction rate theory and partition functions, correlated viscosity [16] as follows:

straight line with slope, $\Delta H^\ddagger / R$ and intercept, $-\Delta S^\ddagger / R$ assuming that ΔH^\ddagger and ΔS^\ddagger be almost independent of temperature. Therefore, ΔH^\ddagger and ΔS^\ddagger can easily be calculated from the slope and intercept of eq. (3) as,

$$\Delta H^\ddagger = \text{Slope} \times R \dots\dots\dots (4)$$

$$\Delta S^\ddagger = -\text{Intercept} \times R \dots\dots\dots (5)$$

The free energy of activation, ΔG^\ddagger , for viscous flow has been calculated by using the simple thermodynamic relation,

$$\Delta G^\ddagger = \Delta H^\ddagger - T\Delta S^\ddagger \dots\dots\dots (6)$$

Calculation of different excess thermodynamic parameters for viscous flow

The excess free energy of activation ($\Delta G^{\ddagger E}$), excess enthalpy of activation ($\Delta H^{\ddagger E}$) and excess entropy of activation ($\Delta S^{\ddagger E}$) for viscous flow were calculated by using the following relations,

$$\Delta G^{\ddagger E} = \Delta G^\ddagger - \Delta G^\ddagger \dots\dots\dots (7)$$

$$\text{or, } \Delta G^{\ddagger E} = \Delta G^\ddagger - (X_1 \Delta G^\ddagger + X_2 \Delta G^\ddagger) \dots\dots\dots (8)$$

$$\eta = \left(\frac{hN}{V_m} \right) \dots\dots\dots$$

V_m

$$\dots\dots\dots \text{). Exp}(\Delta G^\ddagger) / RT \dots\dots\dots (1)$$

$$\Delta H^{\ddagger E} = \Delta H^\ddagger - \Delta H^\ddagger \dots\dots\dots (9)$$

$$\text{or, } \Delta H^{\ddagger E} = \Delta H^\ddagger - (X_1 \Delta H^\ddagger + X_2 \Delta H^\ddagger) \dots\dots\dots (10)$$

Where, ΔG^\ddagger is the free energy of activation per mole for viscous flow, h the Planck's constant (= 6.6262 x

$$\text{And } \Delta S^{\ddagger E} = \Delta S^\ddagger - \Delta S^\ddagger \dots\dots\dots (11)$$

$$\text{or, } \Delta S^{\ddagger E} = \Delta S^\ddagger - (X_1 \Delta S^\ddagger + X_2 \Delta S^\ddagger) \dots\dots\dots (12)$$

10 J.sec), N the Avogadro number (= 6.023 x 10

mol⁻¹), R the molar gas constant (= 8.3145 JK⁻¹mol⁻¹) T the absolute temperature in Kelvin scale and η is the observed viscosity in mPa.s. According to the definition of ΔG^\ddagger eq. (1) reduces to

$$\eta = (hN/V_m) \cdot \exp(-T\Delta S^\ddagger) / RT \dots\dots\dots (2)$$

Where, the subscripts 1 and 2 represent the pure components of the mixture.

The experimentally obtained values of excess properties for a system were fitted by the least square method using Redlich-Kister Eq. (13) of the form:

$$\text{or, } \ln\left(\frac{\eta}{V_m}\right) = - \dots\dots\dots$$

hN

$$\frac{\Delta S^\ddagger}{R} + \Delta H^\ddagger / RT \dots\dots\dots (3)$$

$$(Y)^E = x_1 x_2 \sum_{i \geq 0} (1 - 2x_1)^i B_i \dots\dots\dots (13)$$

Where, ΔH^\ddagger is the enthalpy of activation per mole and ΔS^\ddagger the entropy of activation per mole for viscous flow.

4. Results and discussion

Density and Viscosity of the binary mixture The density and viscosity values of binary mixture as a function of the mole fraction of 2-Bu-OH (x_1) respectively at temperatures (298.15 to 323.15) K can be known by Figure (1-2). The plotting displays the trend of changing these properties. The changes of the properties are evaluated for the changes of each proportion of the mixture component along with the variation of temperatures.




Figure 1: Density trend analysis of the binary mixture




Figure 2: Viscosity trend analysis of the binary mixture

From Figure 2 it can be observed that viscosity of the binary mixture of 2-Bu-OH + m-Xln increases in a very disciplined way with the increasing proportion of 2-Bu-OH but decreases almost linearly with temperature rises. Here the nonpolar m-Xln is a low viscous liquid and when it interacts with the polar one it breaks the intermolecular network of the 2-Bu-OH which was built due to a lot of H-bond as

well as polar-polar interaction. Thus, there was an induction of polarity occurred in m-Xln and became an induced dipole and attached to the polar 2-Bu-OH. Thus, the liquid mixture becomes more viscous. From Figure 2 it can be seen that the rising of viscosity first increases slowly but when the polar proportion is higher it rises sharply upward

Thermodynamic properties measurement

The three types of very important thermodynamic properties viz. entropy, enthalpy and free energy

were very intensively studied for the binary mixture of polar + nonpolar compound. These excess values were also calculated and fitted in Redlich-Kister fitting equation. Figure 3 shows the plot of enthalpy of activation ΔH^\ddagger for the transport process vs. mole fraction of alcohol for the system over the whole range of composition (0 to 1). It is noticed that for the binary system ΔH^\ddagger value (Table 1) continues to increase on addition of alcohol and eventually reach the value of pure 2-Bu-OH because H-bonded liquids, like 2-Bu-OH always require higher enthalpy for activation than that of the other low polar or nonpolar liquids. The excess enthalpies of the binary mixtures of aromatic hydrocarbons with 2-Bu-OH are graphically represented by Figure 4. The excess enthalpy of activation for viscous flow ($\Delta H^{\ddagger E}$) shows the negative values extended over a considerable region of concentrations.

Table 1: Experimental data for enthalpy of activation of the binary mixture

Mole fraction of		Enthalpy (Expt.)	Enthalpy (Theor.)	Excess Enthalpy	Fitting Value
2-Bu-OH (x_1)	m-Xln (x_2)	ΔH^\ddagger kJmol ⁻¹	ΔH^\ddagger_{id} kJmol ⁻¹	$\Delta H^{\ddagger E}$ kJmol ⁻¹	$\Delta H^{\ddagger E*}$ kJmol ⁻¹
0.0000	1.0000	8.2473	8.2473	0.0000	0.0000
0.0500	0.9500	8.8447	9.1483	-0.3036	-0.4519
0.1007	0.8994	8.7808	10.0609	-1.2802	-0.9986
0.1499	0.8501	9.4970	10.9482	-1.4512	-1.5620
0.1998	0.8002	9.8594	11.8470	-1.9876	-2.1274
0.2496	0.7504	10.1993	12.7449	-2.5456	-2.6618
0.2998	0.7002	10.1118	13.6486	-3.5367	-3.1529
0.3498	0.6502	11.0383	14.5502	-3.5119	-3.5863
0.3996	0.6004	11.5978	15.4470	-3.8493	-3.9553
0.4498	0.5502	12.1391	16.3519	-4.2128	-4.2609

0.4993	0.5007	12.7254	17.2432	-4.5178	-4.4925
0.5490	0.4510	13.4728	18.1394	-4.6666	-4.6509
0.5997	0.4003	14.3952	19.0534	-4.6582	-4.7280
0.6493	0.3507	15.0475	19.9470	-4.8995	-4.7104
0.7026	0.2974	16.3868	20.9079	-4.5211	-4.5723
0.7497	0.2503	17.4724	21.7567	-4.2843	-4.3289
0.8001	0.1999	18.9108	22.6649	-3.7542	-3.9175
0.8483	0.1518	19.9537	23.5317	-3.5780	-3.3499
0.9009	0.0991	22.0578	24.4800	-2.4222	-2.4915
0.9503	0.0497	23.9739	25.3712	-1.3973	-1.4104
1.0000	0.0000	26.2660	26.2660	0.0000	0.0000

Enthalpy, Excess Enthalpy and Fitting value of Excess enthalpy of activation for different molar ratios are listed in the table.




Figure3: Enthalpy curve for different mole fractions




Figure 4: Excess enthalpy curve for different mole fractions

Figure 5 is for the plots of entropy of activation ΔS^\ddagger for the viscous flow vs. mole fraction of alcohol for this binary system. The corresponding data are listed in Table 2. The typical nature of the small and negative (ΔS^\ddagger) value of alcohol indicates that during the viscous flow of initial H-bonded order breaks down to form the activated complex of little ordered structure. On the other hand, negative (ΔS^\ddagger) value for the aromatic hydrocarbons revealed that during the flow process the activated complex formed, are much more ordered.

Figure 5 also reveals that with increase of concentration of alcohol, the negative value decreases for the binary system of 2-Bu-OH + m-Xln respectively. It indicates that during the flow process at the certain composition of mixtures, the molecular order of the activated and inactivated state for each of the mixtures is same. The excess entropy $\Delta S^{\#E}$ of activation for the system are shown (Figure 6) as a function of mole fraction of 2-Bu-OH. The plots of $\Delta S^{\#E}$ vs. mole fraction of 2-Bu-OH Figure 6 for the binary mixture show an almost similar trend at those of the excess enthalpy ($\Delta H^{\#E}$). The curve has been found to be concave in nature. The negative excess entropy signifies that the species formed in the activated state are more ordered than what is to be expected to the additive law.

Table 2: Experimental data for Entropy of activation of the binary mixture

Mole fraction of		Entropy (Expt.)	Entropy (Theor.)	Excess Entropy $\Delta S^{#E}$	Fitting Value $\Delta S^{#E*}$
2-Bu-OH (x_1)	m-Xln (x_2)	$\Delta S^{#}$ Jmol ⁻¹ K ⁻¹	$\Delta S^{#}_{id}$ Jmol ⁻¹ K ⁻¹	Jmol ⁻¹ K ⁻¹	Jmol ⁻¹ K ⁻¹
0.0000	1.0000	-53.7654	-53.7654	0.0000	0.0000
0.0500	0.9500	-51.7552	-51.3158	-0.4394	-0.9349
0.1007	0.8994	-71.7320	-48.8346	-3.2086	-2.2531
0.1499	0.8501	-49.7181	-46.4224	-3.2957	-3.6748
0.1998	0.8002	-48.6127	-43.9789	-4.6339	-5.1185
0.2496	0.7504	-47.6356	-41.5379	-6.0977	-6.4842
0.2998	0.7002	-48.1038	-39.0810	-9.0228	-7.7373
0.3498	0.6502	-45.2342	-36.6297	-8.6046	-8.8437
0.3996	0.6004	-43.6499	-34.1915	-9.4584	-9.7901
0.4498	0.5502	-42.1572	-31.7316	-10.4256	-10.5802
0.4993	0.5008	-40.5356	-29.3084	-11.2272	-11.1861
0.5490	0.4510	-38.5053	-26.8720	-11.6333	-11.6068
0.5997	0.4003	-35.9968	-24.3870	-11.6098	-11.8182
0.6493	0.3507	-34.4003	-21.9576	-12.4427	-11.7829
0.7026	0.2974	-30.6157	-19.3452	-11.2704	-11.4360
0.7497	0.2503	-27.6821	-17.0375	-10.6446	-10.8196
0.8001	0.1999	-23.8235	-14.5684	-9.2550	-9.7820
0.8483	0.1518	-21.3038	-12.2120	-9.0918	-8.3609
0.9009	0.0991	-15.6459	-9.6339	-6.0120	-6.2240
0.9503	0.0497	-10.7034	-7.2109	-3.4924	-3.5347
1.0000	0.0000	-4.7782	-4.7782	0.0000	0.0000

Entropy, Excess entropy and Fitting value of excess entropy of activation for different molar ratios are listed in the table.




Figure 5: Entropy curve for different mole fractions




Figure 6: Excess entropy curve for different mole fractions

Table (3-8) shows the variation of free energy of activation ΔG^\ddagger for the viscous flow of the system as a function of 2-Bu-OH under the whole range of composition at temperature 298.15 K to 323.15 K and trend of changing this thermodynamic property

negative, but with the rise of temperature, the values are less negative, i.e., $(\delta\Delta G^\ddagger/\delta T)_p$ is positive. The general nature of the curves does not virtually change with the variation of temperature. The negative excess free energies indicate according to the Eyring

are given by the Figure 7. The ΔG^\ddagger values increase very slowly in the initial stage which is followed by a

$$\ln(\eta V_m) = \frac{\Delta H^\ddagger}{RT} - \Delta S^\ddagger$$

$$-$$

$$\Delta S^\ddagger$$

that the viscous flow

$$= \frac{\Delta H^\ddagger}{RT} - \Delta S^\ddagger$$

relatively greater rise with increasing concentration of 2-Bu-OH. The curves for ΔG^\ddagger for the system are found to be smooth and similar. But a crossover of curves between temperature 298.15 K and 323.15 K is noticed in the system at higher mole fraction of 2-Bu-OH. It is also revealed from the Figure 7 that the nonpolar (m-Xln) rich region the ΔG^\ddagger values increase slightly with the rise of temperature but at the alcohol rich region it is vice versa.

Figure 8 shows the variation of excess free energy $\Delta G^{\ddagger E}$ of activation at 298.15 to 323.15 K over the whole composition range for this binary system (Table 3-8). In each case, the $\Delta G^{\ddagger E}$ values are

$$\Delta G^{\ddagger E} = \Delta G^\ddagger - \Delta G^\ddagger_{ideal}$$

of the solutions of the aromatic hydrocarbons in 2-Bu-OH is enhanced, causing the viscosity to decrease from the values expected ideally.

The free energy of activation is often regarded to be an energy barrier that a molecule must surmount in order to make a hole which is a necessary requirement for a molecule to flow through. The excess values throughout the whole concentration range are negative with minima falling around 0.6 mole fraction of 2-Bu-OH. The negative excess free energy indicates the reduction of energy barrier height and hence increases of viscous flow.

Table3: Experimental data for Free energy of activation of the binary mixture at 298.15 K

Mole fraction of		Free Energy (Expt.) $\Delta G^{\#}$ kJmol^{-1}	Free Energy (Theor.) $\Delta G^{\#}_{id}$ kJmol^{-1}	Excess Free Energy $\Delta G^{\#E}$ kJmol^{-1}	Fitting Value $\Delta G^{\#E*}$ kJmol^{-1}
2-Bu-OH(x_1)	m-Xln(x_2)				
0.0000	1.0000	24.2774	24.2774	0.0000	0.0000
0.0500	0.9500	24.2755	24.4481	-0.1726	-0.1732
0.1007	0.8994	24.2974	24.6210	-0.3235	-0.3268
0.1499	0.8501	24.3205	24.7890	-0.4686	-0.4664
0.1998	0.8002	24.3533	24.9593	-0.6060	-0.6013
0.2496	0.7504	24.4018	25.1294	-0.7276	-0.7286
0.2998	0.7002	24.4540	25.3006	-0.8466	-0.8460
0.3498	0.6502	24.5249	25.4714	-0.9465	-0.9495

0.3996	0.6004	24.6120	25.6412	-1.0293	-1.0364
0.4498	0.5502	24.7083	25.8127	-1.1044	-1.1064
0.4993	0.5008	24.8111	25.9815	-1.1704	-1.1573
0.5490	0.4510	24.9531	26.1512	-1.1981	-1.1903
0.5997	0.4003	25.1277	26.3244	-1.1967	-1.2044
0.6493	0.3507	25.3040	26.4937	-1.1897	-1.1974
0.7026	0.2974	25.5149	26.6757	-1.1608	-1.1627
0.7497	0.2503	25.7259	26.8365	-1.1106	-1.1031
0.8001	0.1999	26.0137	27.0085	-0.9948	-1.0010
0.8483	0.1518	26.3054	27.1727	-0.8673	-0.8571
0.9009	0.0991	26.7226	27.3523	-0.6297	-0.6358
0.9503	0.0497	27.1651	27.5212	-0.3560	-0.3565
1.0000	0.0000	27.6907	27.6907	0.0000	0.0000

Free energy, Excess free energy, and Fitting value of excess free energy for different molar ratios at 298.15 K are listed in the table.

Table4: Experimental data for Free energy of activation of the binary mixture at 303.15 K

Molefractionof		Free Energy (Expt.) $\Delta G^{\#}$ kJmol ⁻¹	FreeEnergy (Theor.) $\Delta G^{\#}_{id}$ kJmol ⁻¹	Excess Free Energy $\Delta G^{\#E}$ kJmol ⁻¹	Fitting Value $\Delta G^{\#E*}$ kJmol ⁻¹
2-Bu-OH(x ₁)	m-Xln(x ₂)				
0.0000	1.0000	24.5462	24.5462	0.0000	0.0000
0.0500	0.9500	24.5343	24.7047	-0.1704	-0.1685
0.1007	0.8994	24.5577	24.8651	-0.3075	-0.3156
0.1499	0.8501	24.5691	25.0212	-0.4521	-0.4480
0.1998	0.8002	24.5963	25.1792	-0.5829	-0.5757
0.2496	0.7504	24.6400	25.3371	-0.6971	-0.6961
0.2998	0.7002	24.6945	25.4960	-0.8015	-0.8074
0.3498	0.6502	24.7511	25.6545	-0.9035	-0.9053
0.3996	0.6004	24.8302	25.8122	-0.9820	-0.9874
0.4498	0.5502	24.9191	25.9713	-1.0523	-1.0535
0.4993	0.5008	25.0138	26.1280	-1.1142	-1.1014

0.5490	0.4510	25.1457	26.2856	-1.1399	-1.1323
0.5997	0.4003	25.3077	26.4463	-1.1387	-1.1453
0.6493	0.3507	25.4760	26.6035	-1.1275	-1.1385
0.7026	0.2974	25.6679	26.7724	-1.1045	-1.1055
0.7497	0.2503	25.8643	26.9217	-1.0574	-1.0490
0.8001	0.1999	26.1328	27.0814	-0.9485	-0.9520
0.8483	0.1518	26.4119	27.2338	-0.8218	-0.8153
0.9009	0.0991	26.8009	27.4005	-0.5997	-0.6046
0.9503	0.0497	27.2187	27.5572	-0.3385	-0.3388
1.0000	0.0000	27.7145	27.7145	0.0000	0.0000

Free energy, Excess free energy, and Fitting value of excess free energy for different molar ratios at 303.15 K are listed in the table.

Table 5: Experimental data of Free energy of activation of the binary mixture at 308.15 K

Molefractionof		Free Energy (Expt.) $\Delta G^{\#}$ kJmol ⁻¹	FreeEnergy (Theor.) $\Delta G^{\#}_{id}$ kJmol ⁻¹	Excess Free Energy $\Delta G^{\#E}$ kJmol ⁻¹	Fitting Value $\Delta G^{\#E*}$ kJmol ⁻¹
2-Bu-OH(x ₁)	m-Xln(x ₂)				
0.0000	1.0000	24.8151	24.8151	0.0000	0.0000

0.0500	0.9500	24.7931	24.9613	-0.1682	-0.1639
0.1007	0.8994	24.8179	25.1093	-0.2914	-0.3043
0.1499	0.8501	24.8177	25.2533	-0.4356	-0.4297
0.1998	0.8002	24.8394	25.3991	-0.5597	-0.5502
0.2496	0.7504	24.8782	25.5448	-0.6666	-0.6637
0.2998	0.7002	24.9350	25.6914	-0.7564	-0.7687
0.3498	0.6502	24.9772	25.8377	-0.8604	-0.8611
0.3996	0.6004	25.0485	25.9832	-0.9347	-0.9385
0.4498	0.5502	25.1298	26.1300	-1.0001	-1.0006
0.4993	0.5008	25.2165	26.2746	-1.0581	-1.0455
0.5490	0.4510	25.3382	26.4200	-1.0818	-1.0743
0.5997	0.4003	25.4877	26.5683	-1.0806	-1.0863
0.6493	0.3507	25.6480	26.7132	-1.0653	-1.0796

0.7026	0.2974	25.8210	26.8691	-1.0481	-1.0483
0.7497	0.2503	26.0027	27.0069	-1.0042	-0.9949
0.8001	0.1999	26.2520	27.1542	-0.9022	-0.9031
0.8483	0.1518	26.5184	27.2948	-0.7764	-0.7735
0.9009	0.0991	26.8791	27.4487	-0.5696	-0.5735
0.9503	0.0497	27.2722	27.5933	-0.3211	-0.3212
1.0000	0.0000	27.7384	27.7384	0.0000	0.0000

Free energy, Excess free energy, and Fitting value of excess free energy for different molar ratios at 308.15 K are listed in the table.

Table 6: Experimental data of Free energy of activation of the binary mixture at 313.15 K

Mole fraction of		Free Energy (Expt.) $\Delta G^\#$ kJmol^{-1}	Free Energy (Theor.) $\Delta G^\#_{id}$ kJmol^{-1}	Excess Free Energy $\Delta G^{\#E}$ kJmol^{-1}	Fitting Value $\Delta G^{\#E*}$ kJmol^{-1}
2-Bu-OH(x_1)	m-Xln(x_2)				
0.0000	1.0000	25.0839	25.0839	0.0000	0.0000
0.0500	0.9500	25.0518	25.2178	-0.1660	-0.1592
0.1007	0.8994	25.0781	25.3535	-0.2754	-0.2930

0.1499	0.8501	25.0663	25.4854	-0.4191	-0.4113
0.1998	0.8002	25.0825	25.6190	-0.5365	-0.5246
0.2496	0.7504	25.1164	25.7525	-0.6361	-0.6313
0.2998	0.7002	25.1755	25.8868	-0.7113	-0.7300
0.3498	0.6502	25.2034	26.0208	-0.8174	-0.8169
0.3996	0.6004	25.2667	26.1541	-0.8874	-0.8896
0.4498	0.5502	25.3406	26.2886	-0.9480	-0.9477
0.4993	0.5008	25.4191	26.4211	-1.0020	-0.9896
0.5490	0.4510	25.5307	26.5543	-1.0236	-1.0162
0.5997	0.4003	25.6676	26.6902	-1.0226	-1.0272
0.6493	0.3507	25.8200	26.8230	-1.0030	-1.0206
0.7026	0.2974	25.9741	26.9659	-0.9918	-0.9912
0.7497	0.2503	26.1411	27.0920	-0.9510	-0.9408

0.8001	0.1999	26.3711	27.2270	-0.8560	-0.8542
0.8483	0.1518	26.6250	27.3559	-0.7309	-0.7317
0.9009	0.0991	26.9573	27.4968	-0.5395	-0.5424
0.9503	0.0497	27.3257	27.6293	-0.3036	-0.3035
1.0000	0.0000	27.7623	27.7623	0.0000	0.0000

Free energy, Excess free energy, and Fitting value of excess free energy for different molar ratios at 313.15 K are listed in the table.

Table 7: Experimental data for Free energy of activation of the binary mixture at 318.15 K

Mole fraction of		Free Energy (Expt.) $\Delta G^{\#}$ kJmol^{-1}	Free Energy (Theor.) $\Delta G^{\#}_{id}$ kJmol^{-1}	Excess Free Energy $\Delta G^{\#E}$ kJmol^{-1}	Fitting Value $\Delta G^{\#E*}$ kJmol^{-1}
2-Bu-OH(x_1)	m-Xln(x_2)				
0.0000	1.0000	25.3527	25.3527	0.0000	0.0000
0.0500	0.9500	25.3106	25.4744	-0.1638	-0.1545
0.1007	0.8994	25.3383	25.5977	-0.2594	-0.2818
0.1499	0.8501	25.3149	25.7175	-0.4026	-0.3929
0.1998	0.8002	25.3255	25.8389	-0.5134	-0.4990
0.2496	0.7504	25.3546	25.9601	-0.6056	-0.5989
0.2998	0.7002	25.4160	26.0822	-0.6661	-0.6913

0.3498	0.6502	25.4296	26.2040	-0.7744	-0.7727
0.3996	0.6004	25.4850	26.3251	-0.8401	-0.8406
0.4498	0.5502	25.5514	26.4473	-0.8959	-0.8948
0.4993	0.5008	25.6218	26.5676	-0.9458	-0.9336
0.5490	0.4510	25.7232	26.6887	-0.9654	-0.9582
0.5997	0.4003	25.8476	26.8121	-0.9645	-0.9681
0.6493	0.3507	25.9920	26.9328	-0.9408	-0.9617
0.7026	0.2974	26.1272	27.0626	-0.9354	-0.9340
0.7497	0.2503	26.2795	27.1772	-0.8977	-0.8867
0.8001	0.1999	26.4902	27.2999	-0.8097	-0.8053
0.8483	0.1518	26.7315	27.4169	-0.6855	-0.6899
0.9009	0.0991	27.0355	27.5450	-0.5095	-0.5113

0.9503	0.0497	27.3792	27.6654	-0.2862	-0.2858
1.0000	0.0000	27.7862	27.7862	0.0000	0.0000

Free energy, Excess free energy, and Fitting value of excess free energy for different molar ratios at 318.15 K are listed in the table.

Table 8: Experimental data for Free energy of activation of the binary mixture at 323.15 K

Mole fraction of		Free Energy (Expt.)	Free Energy (Theor.)	Excess Free Energy	Fitting Value
2-Bu-OH(x_1)	m-Xln(x_2)	$\Delta G^{\#}$ kJmol ⁻¹	$\Delta G^{\#}_{id}$ kJmol ⁻¹	$\Delta G^{\#E}$ kJmol ⁻¹	$\Delta G^{\#E*}$ kJmol ⁻¹
0.0000	1.0000	25.6216	25.6216	0.0000	0.0000
0.0500	0.9500	25.5694	25.7310	-0.1616	-0.1498
0.1007	0.8994	25.5985	25.8418	-0.2433	-0.2705
0.1499	0.8501	25.5634	25.9496	-0.3862	-0.3745
0.1998	0.8002	25.5686	26.0588	-0.4902	-0.4734
0.2496	0.7504	25.5927	26.1678	-0.5751	-0.5665
0.2998	0.7002	25.6566	26.2776	-0.6210	-0.6526
0.3498	0.6502	25.6557	26.3871	-0.7314	-0.7284
0.3996	0.6004	25.7032	26.4960	-0.7928	-0.7917
0.4498	0.5502	25.7622	26.6059	-0.8438	-0.8419
0.4993	0.5008	25.8245	26.7142	-0.8897	-0.8777

0.5490	0.4510	25.9158	26.8230	-0.9073	-0.9002
0.5997	0.4003	26.0276	26.9341	-0.9065	-0.9090
0.6493	0.3507	26.1640	27.0426	-0.8786	-0.9028
0.7026	0.2974	26.2803	27.1593	-0.8791	-0.8768
0.7497	0.2503	26.4179	27.2624	-0.8445	-0.8326
0.8001	0.1999	26.6093	27.3727	-0.7634	-0.7564
0.8483	0.1518	26.8380	27.4780	-0.6400	-0.6480
0.9009	0.0991	27.1138	27.5932	-0.4794	-0.4802
0.9503	0.0497	27.4327	27.7014	-0.2687	-0.2681
1.0000	0.0000	27.8101	27.8101	0.0000	0.0000

Free energy, Excess free energy, and Fitting value of excess free energy for different molar ratios at 323.15 K are listed in the table.




Figure 7: Free energy curves for different mole fractions at different temperatures




Figure 8: Excess free energy curves for different mole fractions at different temperatures

5. Conclusion

Studies on the thermodynamic properties along with their respective excess values in liquids and liquid mixtures is essential to understand the molecular interactions between unlike molecules, to develop new theoretical models [17] as well as engineering applications in pharmaceutical process industries. An understanding of the thermodynamic properties of 2-Bu-OH and m-Xln and their binary system have been measured at temperatures (298.15 to 323.15) K and in atmospheric pressure. Excess thermodynamic properties of binary mixture solvents were calculated and fitted with the Redlich–Kister equation. Thermodynamic properties derived from viscosity and its related parameters are also important in designing industrial equipments with better precision [18]. In these mixtures, the predominance of physical effect may arise due to dipole-induced dipole interaction resulting in disruption in the favorable orientation due to breaking of self-network built by H-bond among the polar molecules and rising dispersion force of nonpolar liquid. The observed excess thermodynamic values in all the mixture

indicate the significant interaction between the unlike molecules.

References

1. The use of organic solvents in the pharmaceutical sector. 1 Grodowska K, Parczewski A. *Acta Poloniae Pharmaceutica et Drogue Research*, 67, 3–12 (2010).
2. Organic solvents for pharmaceutical parenterals and embolic liquids: a review of toxicity data. Mottu F, Laurent A, Rufenacht DA, Doelker E. To cite this article: *PDA J Pharm Sci Technol* 54 (2000): 456-69.
3. American Chemistry Council, "Solvents for Pharmaceutical Manufacturing," 2016.
4. Residual solvents analysis (volatile organic compounds; OVI) by Intertec (2016).
5. Yadav, R., Yadav, N., & Kharya, M.D.

- Quality assurance for medicines containing trace amounts of solvents. The full citation for this article is W J Pharmacy Pharma Sci 3 (2014): 526-538.
6. Residual solvents, USP 37, Chemical test 467:1-37.
 7. Seventh, Industry Guidance Q3C Tables and List,
 8. Health and Human Services (U.S.). (2016).
 9. Frank, Howard, Thomas WM, Fookson A, Pomerantz P, Brooks DB. Preparation and physical characteristics of various aliphatic hydrocarbons and intermediates. There was a report on this in J. Res.
 10. For more information on the dangers of 2-butanol in the workplace, see NIOSH (2016). Xylene, Pennsylvania Department of Health (2016).
 11. Uniform growth and simplified production are the results of Rahman et al.'s methodical synthesis of carbon nanotubes from a mixture of aliphatic and aromatic compounds (11). 2 (2020): 001-009 Journal of Nanotechnology Research
 12. Study of acoustic characteristics of binary mixes of a nonpolar liquid and a polar liquid at various frequencies.